

Synthesis and characterization of templated pentaborate(1-) salts: X-ray structure of [(2-HOCH₂CH₂)C₄H₇NMeH][B₅O₆(OH)₄]-0.3H₂O

Jones, Charlotte; Beckett, Michael; Coles, Simon J.; Davies, Richard; Horton, Peter N.

Phosphorus, Sulfur and Silicon and the Related Elements

DOI:

[10.1080/10426507.2015.1128921](https://doi.org/10.1080/10426507.2015.1128921)

Published: 01/08/2016

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Jones, C., Beckett, M., Coles, S. J., Davies, R., & Horton, P. N. (2016). Synthesis and characterization of templated pentaborate(1-) salts: X-ray structure of [(2-HOCH₂CH₂)C₄H₇NMeH][B₅O₆(OH)₄]-0.3H₂O. *Phosphorus, Sulfur and Silicon and the Related Elements*, 191(4), 628-630. <https://doi.org/10.1080/10426507.2015.1128921>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Synthesis and characterization of templated pentaborate(1-) salts: X-ray structure of $[(2\text{-HOCH}_2\text{CH}_2)\text{C}_4\text{H}_7\text{NMeH}][\text{B}_5\text{O}_6(\text{OH})_4] \cdot 0.3\text{H}_2\text{O}$

Charlotte L. Jones,¹ Michael A. Beckett*,¹ Simon J. Coles,² R. Andrew Davies,¹ Peter N. Horton.²

¹School of Chemistry, Bangor University, Bangor, Gwynedd, LL57 2UW, UK

²School of Chemistry, University of Southampton, Southampton, SO17 1BJ, UK

GRAPHICAL ABSTRACT



Abstract The synthesis of pyrrolidinium pentaborate(1-) and four substituted pyrrolidinium pentaborate(1-) salts from $\text{B}(\text{OH})_3$ in aqueous solution are reported. Characterization (IR, NMR, single-crystal XRD) of $[(2\text{-HOCH}_2\text{CH}_2)\text{C}_4\text{H}_7\text{NH}_2][\text{B}_5\text{O}_6(\text{OH})_4] \cdot 0.3\text{H}_2\text{O}$ is described in detail. The cation is the largest of the five pyrrolidinium cations studied. A consequence of this is that its solid-state structure has anion-anion interactions ($\alpha\alpha\alpha\gamma$) which are different from the other four ($\alpha\alpha\alpha\beta$) structures. There are also stabilizing cation-anion H-bond interactions.

Keywords Pentaborate(1-); H-bonds; Pyrrolidinium; X-ray structure; Oxidoborate

Received xx yyyy 2015; accepted xx yyyy 2015

We thank the EPSRC for use of the NCS Service

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423523. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

INTRODUCTION

Pentaborate(1-) salts, [NMC][B₅O₆(OH)₄], are often prepared from the reaction of organic bases with B(OH)₃ in aqueous solution, where the protonated organic base is a non-metal cation (NMC).¹ Occasionally, salts containing rarer anions e.g. three to fifteen B atoms, have also been obtained.²⁻⁹ In basic aqueous solutions, B(OH)₃ forms a dynamic combinatorial library (DCL) of polyborate anions whose concentrations are pH and boron concentration dependent.¹⁰⁻¹¹ Recently, we noted from DFT calculations (gas-phase)¹² on the relative stabilities of the polyborate anions that the pentaborate(1-) anion is not the most stable and yet despite this, [NMC][B₅O₆(OH)₄] salts are readily formed. Strong anion-anion H-bond interactions are present in [NMC][B₅O₆(OH)₄] salts and these we believe are responsible for their facile formation. Herein, we report the synthesis of several pentaborate(1-) salts partnered with substituted pyrrolidinium cations, explore steric effects of the cation, and describe in detail the synthesis of the salt [(2-HOCH₂CH₂)C₄H₇NMeH][B₅O₆(OH)₄] (**1**).

RESULTS AND DISCUSSION

Synthesis and characterization


The pyrrolidinium pentaborate salts [(2-HOCH₂CH₂)C₄H₇NMeH][B₅O₆(OH)₄]0.3H₂O (**1**), [(2-HOCH₂)C₄H₇NH₂][B₅O₆(OH)₄]0.5H₂O (**2**), [C₄H₈NMeH][B₅O₆(OH)₄]0.5CH₃COCH₃ (**3**), [C₄H₈NMe₂][B₅O₆(OH)₄] (**4**), and [C₄H₈NH₂][B₅O₆(OH)₄] (**5**), were all prepared in high yields from the reaction of the free base (**1-3**, **5**), or quaternary amine hydroxide salt (**4**), with B(OH)₃ in a 1:5 molar ratio. Characterization data including single-crystal XRD structural analyses for salts **2-5** have been recently been reported elsewhere.¹³

Characterization data¹⁴ (IR, NMR) obtained for **1** were in agreement with those previously reported for related pyrrolidinium and other non-metal cation pentaborate salts. ¹H and ¹³C NMR spectra (obtained in D₂O) were fully consistent with those expected for the cation, with the NH, OH and BOH protons overlapping and represented by a broad singlet at ~4.7 ppm due to rapid exchange. Three characteristic signals at ~17, 13 and 1 ppm were observed in ¹¹B NMR spectra, obtained in concentrated aqueous solutions (D₂O), which are assigned to B(OH)₃/[B(OH)₄]⁻, [B₃O₃(OH)₄]⁻ and the 4-coordinate centre of [B₅O₆(OH)₄]⁻, respectively.¹⁵ These species are

observed due to the complex borate equilibria observed in aqueous solution.^{11,16} The IR spectrum of **1** clearly shows the diagnostic band¹⁶ of pentaborate salts at $\sim 925\text{ cm}^{-1}$.

X-ray structure of $[(2\text{-HOCH}_2\text{CH}_2)_4\text{C}_4\text{H}_7\text{NMeH}][\text{B}_5\text{O}_6(\text{OH})_4]$

The crystallographic data for compound **1** is given in a footnote¹⁷ with full details given in the supplementary data.¹⁸ The ionic salt is comprised of the expected (2-hydroxyethyl)-*N*-methylpyrrolidinium(1+) cation, partnered with a pentaborate(1-) anion, as shown in Figure 1. The cation was found to be disordered over multiple sites with three significant positions of occupancy identified (4:3:3 ratio) and one of which also included a solvated water molecule. All the disordered versions of the pyrrolidinium ring have an envelope conformation with both substituents equatorial. The bond lengths and internuclear angles observed within the boroxyl (B_3O_3) rings of the pentaborate anions of **1** are within the ranges observed for previously reported $[\text{NMC}][\text{B}_5\text{O}_6(\text{OH})_4]$ structures.^{1,6,15,16} The bond lengths and internuclear angles also lie within the ranges found in related boroxole (B_3O_3) structures containing both 3- and 4-coordinate B centres bound to O.¹⁹



Received xx yyyy 2015; accepted xx yyyy 2015

We thank the EPSRC for use of the NCS Service

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423523. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Figure 1 The structure of $[(2\text{-HOCH}_2\text{CH}_2)\text{-C}_4\text{H}_7\text{NMeH}][\text{B}_5\text{O}_6(\text{OH})_4]$ (**1**), showing the atomic numbering scheme.

Structure **1** possesses a giant H-bonded anionic lattice, with the substituted pyrrolidinium cations (and the H_2O) sitting within the cavities of the lattice. The anions arrange themselves in a series of ‘planes’ (Figure 2) H-bonded together *via* $\text{R}_2^2(8)$ reciprocal pair interactions²¹ to $\alpha\alpha\gamma$ acceptor sites, and the ‘planes’ are linked through further reciprocal- α $\text{R}_2^2(8)$ interactions.




Figure 2 Anionic lattice of **1** viewed along the a axis; the $[(2\text{-HOCH}_2\text{CH}_2)\text{C}_4\text{H}_7\text{NMeH}]^+$ cations sit H-bonded within the cavities of the lattice. The $\text{R}_2^2(8)$ reciprocal- α H-bond interactions are shown in blue.

The anions of the H-bonded lattice are in a different arrangement to that observed for **3-5** that have a familiar $\alpha\alpha\alpha\beta$ ‘brickwall’ structure.^{1,15} It is interesting to compare the structures of compounds **1** with **4** and **5**. The brickwall structure is sufficiently flexible to accommodate cations of variable sizes (within limits). *e.g.* the N,N -dimethylated pyrrolidinium cation in **4** has the same structure as the unsubstituted cation in **5**, despite a formula unit increase in volume of 13.3%.¹³ The formula unit volume of **1** has increased by a further 10.3% compared to **4**, as a consequence of the larger substituted cation. This, we believe, is responsible for switching the structure from the brickwall to the $\alpha\alpha\alpha\gamma$ arrangement. In this structure three strong reciprocal-

Received xx yyyy 2015; accepted xx yyyy 2015

We thank the EPSRC for use of the NCS Service

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423523. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

α bonds (-21 kJ mol^{-1}) are maintained and the H-bond energy for a reciprocal- γ interaction is the same as that of a C(8) β -chain (-16 kJ mol^{-1}).¹³ The cation in **1** is also ‘non-innocent’ and cation-anion H-bond interactions *e.g.* N11H11...O4’, N11H11...O9’, and O11H11B...O8’, further stabilize the self-assembled crystal structure. Full details are available in the supplementary information.

SUMMARY

The synthesis and crystal structures of pyrrolidinium pentaborate(1-) and 4 substituted pyrrolidinium pentaborates are reported. The larger cation present in **1** is unable to fit into the brickwall structure and as a result **1** adopts a similar but different anionic lattice of comparable energy. The solid-state structure of **1** is further stabilized by cation-anion H-bond interactions.

REFERENCES

1. (a) Freyhardt, C.C.; Felsche, J.; Engelhardt, G.; *Z. Naturforsch. B: Chem. Sci.*, **1993**, 48, 978-985. (b) Freyhardt, C.C.; Wiebcke, M.; Felsche, J.; Engelhardt, G.; *J. Inclusion Phenom. Mol. Recognit. Chem.*, **1994**, 18, 161-175.
2. (a) Schubert, D.M.; Visi, M.Z.; Knobler, C.B. *Inorg. Chem.* **2008**, 47, 2017-2023; (b) Beckett, M.A.; Horton, P.N.; Hursthouse, M.B.; Timmis, J.L. *RSC Adv.* **2013**, 3, 15185-15191.
3. (a) Wang, G.-M.; Sun Y.-Q.; Yang, G.-Y. *J. Solid State Chem.* 2004, 177, 4648-4654; (b) Weakley, T.J.R.; *Acta Crystallogr. Sect. C: Cryst. Struct. Commun.*, **1985**, C41, 377-379; (c) Beckett, M.A.; Horton, P.N.; Coles, S.J.; Martin, D.W.; *Inorg. Chem.*, **2011**, 50, 12215-12218.
4. Liu, M.-C.; Zhou, P.; Yao, H.-G.; Ji, S.-H.; Zhang, R.-C.; Ji, M.; An, Y.-L.; *Eur. J. Inorg. Chem.*, **2009**, 4622-4624.
5. (a) Schubert, D.M.; Visi, M.Z.; Khan, S.; Knobler, C.B.; *Inorg. Chem.*, **2008**, 47, 4740-4745; (b) Beckett, M.A.; Horton, P.N.; Hursthouse, M.B.; Timmis, J.L.; Varma, K.S.; *Dalton Trans.*, **2012**, 41, 4396-4403.
6. Visi, M.Z.; Knobler, C.B.; Owen, J.J.; Khan, M.I.; Schubert, D.M.; *Cryst. Growth Des.*, **2006**, 6, 538-545.
7. (a) Schubert, D.M.; Smith, R.A.; Visi, M.Z.; *Glass Technol.*, **2003**, 44, 63-70; (b) Schubert, D.M.; Visi, M. Z.; Knobler, C.B.; *Inorg. Chem.*, **2000**, 39, 2250-2251.
8. Liu, Z.H.; Li, L.Q.; Zhang, W.J.; *Inorg. Chem.*, **2006**, 45, 1430-1432.
9. Merlino, S.; Sartori, F.; *Science*, **1971**, 171, 377-379.
10. Corbett, P.T.; Leclaire, J.; Vial, L.; West, K.R.; Wietor, J.-L.; Sanders, J.K.M.; Otto, S.; *Chem. Rev.*, **2006**, 106, 3652-3711.
11. (a) Farmer, J.B.; *Adv. Inorg. Chem.*, **1982**, 25, 187-237; (b) Salentine, C.G.; *Inorg. Chem.*, **1983**, 22, 3920-3924.

Received xx yyyy 2015; accepted xx yyyy 2015

We thank the EPSRC for use of the NCS Service

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423523. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

12. Beckett, M.A.; Davies, R.A.; Thomas, C.D.; *Comput. Theor. Chem.*, **2014**, 1044, 74–79.
13. Beckett, M.A.; Coles, S.J.; Davies, R.A.; Horton, P.N.; Jones, C.L. *Dalton Trans.* **2015**, 44, 7032–7040.
14. Characterization data for **1**: C₇H_{20.60}B₅NO_{11.30}. Elem. Anal Calc: C, 23.8; H 5.9; N, 4.0. Found: C, 23.8; H 5.7; N, 3.9%. Mp >300 °C. IR (KBr/cm⁻¹): 3399(br,s), 2973(m), 2893(m), 1639(m), 1432(m), 1315(s), 1100(s), 1027(s), 923(vs), 776(s) 707(s). NMR (400 MHz, D₂O) δ¹H/ppm: 3.82–3.76 1H (m), 3.72–3.61 2H (m), 3.46–3.38 1H (m), 3.18–3.12 1H (q, 8 Hz) 2.90 3H (s), 2.41–2.32 1H (m), 2.19–2.01 3H (m), 1.86–1.76 2H (m). δ¹¹B/ppm: 16.8, 13.2, 1.2. δ¹³C/ppm: 66.8 (CH), 58.2 (CH₂), 55.9 (CH₂), 38.9 (CH₃), 32.0 (CH₂), 28.8 (CH₂), 21.1 (CH₂).
15. Beckett, M.A.; Horton, P.N.; Hursthouse, M.B.; Knox, D.A.; Timmis, J.L.; *Dalton Trans.*, **2010**, 39, 3944–3951.
16. (a) Beckett, M.A.; Horton, P.N.; Coles, S.J.; Kose, D.A.; Kreuziger, A.-M.; *Polyhedron*, **2012**, 38, 157–161; (b) Beckett, M.A.; Horton, P.N.; Hursthouse, M.B.; Timmis, J.L.; Varma, K.S.; *Collect. Czech. Chem. Commun.*, **2010**, 75, 971–980.
17. A Rigaku AFC12 goniometer equipped with an enhanced sensitivity (HG) Saturn724+ detector mounted at the window of an FR-E+ SuperBright molybdenum rotating anode generator with VHF Varimax optics (70μm focus) was used. X-ray data for **1**: C₇H_{20.60}B₅NO_{11.30}, M = 353.69; monoclinic space group P2₁/c, a = 10.0014(7) Å, b = 11.4193(8) Å, c = 14.8649(10) Å, α = 90°, β = 104.6400(10)°, γ = 90°; U = 1642.6(2) Å³; Z = 4; T = 100(2) K; Wavelength = 0.71075 Å; density (calc) = 1.430 Mg/m³; Absorption coefficient 0.125 mm⁻¹; Crystal: colourless prism, 0.110 x 0.100 x 0.080 mm³; F(000) 740; θ range for data collection 3.073 – 27.484°; Index ranges –12 ≤ h ≤ 12, –14 ≤ k ≤ 14, –19 ≤ l ≤ 18; 22146 reflections collected with 3746 independent reflections [R_{int} = 0.0321]; Completeness to θ = 25.242°: 99.7%; absorption correction: semi-empirical from equivalents; refinement method: full-matrix least squares on F²; data/restraints/parameters 3746/244/261; goodness-of-fit on F², 1.103; Final R indices [F² > 2σ(F²)] R1 = 0.0787, wR2 = 0.2322; R indices (all data) R1 = 0.0944, wR2 = 0.2484; extinction coefficient n/a; largest diff. peak and hole 0.743 and -0.509 e Å⁻³.
18. (a) Coles, S.J.; Gale, P.A. *Chem. Sci.* **2012**, 3, 683–689; (b) Rigaku, CrystalClear-SM Expert 3.1 **2013**, b27; (c) Sheldrick, G.M. *Acta Crystallogr.* **2015**, C71, 3; (d) Farrugia, L.J.; *J. Appl. Crystallogr.* **1997**, 30, 565; (e) Palatinus, L.; Chapuis, G.; *J. Appl. Crystallogr.* **2007**, 40, 786–790.
19. (a) Beckett, M.A.; Coles, S.J.; Light, M.E.; Fischer, L.; Stiefvater-Thomas, B.M.; Varma, K.S. *Polyhedron* **2006**, 25, 1011–1016; (b) Beckett, M.A.; Hibbs, D.E.; Hursthouse, M.B.; Owen, P.; Malik, K.M.A.; Varma, K.S. *Main Group Chem.* **1998**, 2, 251–258; (c) Beckett, M.A.; Strickland, G.C.; Varma, K.S.; Hibbs, D.E.; Hursthouse, M.B.; Malik, K.M.A. *J. Organomet. Chem.* **1997**, 535, 33–41.
20. Etter, M.C.; *Acc. Chem. Res.*, **1990**, 23, 120–126.

Received xx yyyy 2015; accepted xx yyyy 2015

We thank the EPSRC for use of the NCS Service

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423523. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.